

Correlations and Critical Behavior in the $SU(2)$ Gluodynamics

V.G. Bornyakov^{1,3,4}, V.V. Bryzgalov¹, V.K. Mitrjushkin², and
R.N. Rogalyov¹

¹ IHEP, Protvino; ² JINR, Dubna, ³ ITEP, Moscow, ⁴ FEFU, Vladivostok

19.06.2018

- ▶ Phase transitions for pedestrains
- ▶ Phase transition in $SU(2)$
- ▶ Correlation between the **asymmetry** and the **Polyakov loop**
- ▶ Correlation between the **longitudinal propagator** and the **Polyakov loop**
- ▶ Regression analysis
- ▶ Evaluation of the critical exponents and amplitudes
- ▶ Conditional distributions of the longitudinal propagator
- ▶ Conclusions

Ising model

$$\sigma_n = \pm 1$$

Finite-volume lattice: $\vec{n} = (n_1, \dots, n_D)$, $1 \leq n_\mu \leq L$, $n_\mu \in \mathbf{N}$

Infinite-volume lattice: $\vec{n} \in \mathbb{Z}^D$,

$$H = -J \sum_{|\vec{i}-\vec{j}|=1} \sigma_{\vec{a}\vec{i}} \sigma_{\vec{a}\vec{j}} - h \sum_{\vec{i} \in \mathbb{Z}^D} \sigma_{\vec{a}\vec{i}}$$

$$Z = \sum_{\sigma_n} e^{-H[\sigma]/T} = \exp\left(-\frac{F}{T}\right)$$

$$m = \frac{\partial F}{\partial h}, \quad S = -\frac{\partial F}{\partial T}, \quad \chi = \frac{\partial^2 F}{\partial h^2}, \quad c = -T \frac{\partial^2 F}{\partial T^2},$$

Phase transition: singular behavior of $F(T)$ at $T = T_c$

Critical exponents

$$\tau = \frac{T - T_c}{T_c}$$

$$m|_{h=0} \simeq C_\beta (-\tau)^\beta \quad (\tau < 0)$$

$$|m|_{\tau=0} \simeq C_\delta |h|^{1/\delta} \quad (\tau = 0)$$


$$\chi = \frac{\partial m}{\partial h} \simeq C_\gamma |\tau|^{-\gamma}$$


$$G(\vec{k})|_{\tau=0} = a^D \sum_{\vec{n} \in \mathbb{Z}^D} \langle \sigma_{a\vec{n}} \sigma_{\vec{0}} \rangle e^{ia\vec{n}\vec{k}} \simeq \frac{C_\eta}{|\vec{k}|^{2-\eta}}$$


correlation length ξ : $\langle \sigma_{\vec{x}} \sigma_{\vec{0}} \rangle|_{\tau \neq 0} \sim e^{-|\vec{x}|/\xi}$


$$\xi \simeq C_\xi |\tau|^{-\nu}$$

heat capacity : $c_{h=0} \simeq C_c |\tau|^{-\alpha}$











Probability density



Probability density



Binder 1981; Mitrjuskin, Zadorozhny 1986

$$(\vec{x}, x_4) \in \Lambda(N_t \times N_s^3), \quad \quad N_t = 8, \quad 32 \leq N_s \leq 88$$

$$L(\vec{x}) = \frac{1}{2} \operatorname{Tr} \prod_{x_4=1}^{N_t} U(\vec{x}, x_4; \mu = 4)$$

$$\mathcal{P} = \frac{1}{N_s^3} \sum_{\vec{x}} L(\vec{x})$$

$$\langle L(\vec{x})L(\vec{0}) \rangle \simeq A \exp \left(- \frac{|\vec{x}|}{\xi} \right), \quad |\vec{x}| \rightarrow \infty$$

$$G(\vec{p}) = \frac{1}{N_s^3} \sum_{\vec{x}} \langle L(\vec{x})L(\vec{0}) \rangle e^{i\vec{p}\vec{x}}$$

Critical exponents and amplitudes

$$\tau = \frac{T - T_c}{T_c}; \quad \tau > 0 - \text{deconfinement}$$

$$\langle \mathcal{P} \rangle \simeq B \tau^\beta$$

$$\xi \simeq \frac{f_\pm}{|\tau|^\nu}$$

$$\langle \mathcal{P}^2 \rangle - \langle \mathcal{P} \rangle^2 = G(\vec{0}) \simeq \frac{C_\pm}{N_s^3 |\tau|^\gamma}$$

$$\tau = 0 : \quad G(\vec{p}) \simeq \frac{H}{|\vec{p}|^{2-\eta}}$$

Critical exponents

and amplitudes

3D Ising

F.Kos, D.Poland et al.
JHEP (2016)

$$\beta = 0.326419(3)$$

$$\gamma = 1.237075(10)$$

$$\eta = 0.036298(2)$$

$$\nu = 0.629971(4)$$

$SU(2), 4D$

J.Engels, T.Schiedeler 1998

$$B = 0.825(1)$$

$$C_+ = 0.0587(8), \quad C_- = 0.01243(12)$$

C_+/C_- is universal; for the 3D Ising universality class

$$C_+/C_- = 4.75(3) \text{ [1998]}$$


$$\langle \alpha(x) \alpha(y) \rangle = \frac{1}{|x-y|^{2\Delta\alpha}}$$

$$\langle A(x) B(y) C(z) \rangle = \frac{f_{ABC}}{|x-y|^{\Delta_A + \Delta_B - \Delta_C} |y-z|^{\Delta_B + \Delta_C - \Delta_A} |x-z|^{\Delta_A + \Delta_C - \Delta_B}}$$

$$\nu = \frac{1}{3 - \Delta_\epsilon}, \quad \gamma = \frac{3 - 2\Delta_6}{3 - \Delta_\epsilon}$$

$$\langle G(x_1) G(x_2) G(x_3) G(x_4) \rangle = \sum_{\sigma} \langle \overbrace{x_1 x_2}^1 \overbrace{x_3 x_4}^4 \rangle_{\sigma} = \sum_{\sigma} f_{GGG}^2 C_{\Delta_6 \Delta_6}^{\Delta_6}(x_1, \dots, x_4)$$

$$\sum_{\sigma} f_{GGG}^2 (C_{\Delta_6 \Delta_6}^{\Delta_6}(x_1, x_2, x_3, x_4) - C_{\Delta_6 \Delta_6}^{\Delta_6}(x_3, x_2, x_1, x_4)) = 0$$



A.Maas, J.Pawlowski, L von Smekal, D.Spielmann, 2011

A.Maas *et al.*, 2011 (6×48^3):

$$M_E(\tau) = m_{gribov} + \theta(\tau)\mathcal{M}_+\tau^{\gamma_+/2} + \theta(-\tau)\mathcal{M}_-\tau^{\gamma_-/2} \quad (1)$$

$$m_{gribov} = 0.25^{+3}_{-2}; \quad \mathcal{M}_+ = 1.5^{+1}_{-3}, \quad \mathcal{M}_- = -0.07^{+736}_{-17}; \quad (2)$$

$$\gamma_+ = 1.54^{+12}_{-0.05}, \quad \gamma_- = 0.6^{+45}_{-5}$$

Our **previous** result 2015, ($N_t = 8$, extrapolation to the infinite-volume limit):

$$m_{gribov} = 0.217(3); \quad \mathcal{M}_+ = 0.93(11), \quad \mathcal{M}_- = -1.23(19); \quad (3)$$

$$\gamma_+ = \gamma_- = 0.63(3)$$

The Chromo-Electric-Magnetic Asymmetry

$$\begin{aligned}\langle A_E^2 \rangle &= g^2 \langle A_4^a(x) A_4^a(x) \rangle, \\ \langle A_M^2 \rangle &= g^2 \langle A_i^a(x) A_i^a(x) \rangle.\end{aligned}\tag{4}$$

The quantity of particular interest is the (color) electric-magnetic asymmetry introduced by Chernodub and Ilgenfritz in 2008:

$$\langle \Delta_{A^2} \rangle \equiv \langle A_E^2 \rangle - \frac{1}{3} \langle A_M^2 \rangle. \tag{5}$$

Later we will use the dimensionless quantity

$$\Delta_{A^2} = \frac{\langle A_E^2 \rangle - \frac{1}{3} \langle A_M^2 \rangle}{T^2}. \tag{6}$$

We work in the Landau gauge $\partial_\mu A_\mu^a = 0$

Definition of the longitudinal (L) and transverse (T) propagators:

$$D_{\mu\nu}^{ab}(p) = \delta_{ab} \left(P_{\mu\nu}^T(p) D_T(p) + P_{\mu\nu}^L(p) D_L(p) \right),$$

where $P_{\mu\nu}^{T;L}(p)$ - orthogonal transverse (longitudinal) projectors

$$D_L(p) = \frac{1}{3} \sum_{a=1}^3 \langle A_0^a(p) A_0^a(-p) \rangle$$

$$D_T(p) = \begin{cases} \frac{1}{6} \sum_{a=1}^3 \sum_{i=1}^3 \langle A_i^a(p) A_i^a(-p) \rangle & p \neq 0 \\ \frac{1}{9} \sum_{a=1}^3 \sum_{i=1}^3 \langle A_i^a(p) A_i^a(-p) \rangle & p = 0 \end{cases}$$

We study critical behavior of the quantities

$$\mathcal{A} = \Delta_{A^2} - \Delta_{A^2}^C \quad (7)$$

and

$$\mathcal{D} = D_L(0) - D_L^C(0) , \quad (8)$$


Asymptotic expansions of Δ_{A^2} and \mathcal{D}

in $\tau = \frac{T - T_c}{T_c}$ at $\tau \rightarrow 0_+$ have the form

$$\mathcal{A} \simeq B_{\mathcal{A}} \tau^{\beta_{\mathcal{A}}} , \quad (9)$$


$$\mathcal{D} \simeq B_{\mathcal{D}} \tau^{\beta_{\mathcal{D}}} , \quad (10)$$

We evaluate the critical exponents $\beta_{\mathcal{A}}$ and $\beta_{\mathcal{D}}$
and amplitudes $B_{\mathcal{A}}$ and $B_{\mathcal{D}}$.





Unjustified assumptions:

- ▶ **Maas' et al.:** Correlator $\langle A_0(\vec{x})A_0(\vec{0}) \rangle$ is associated with the same critical exponent (γ) as that of Polyakov loops
- ▶ **Our :** Negative Polyakov-loop sector can be safely ignored




Chernodub, Ilgenfritz 2008







Average value of Δ_{A^2} versus T , $T_c \rightarrow \beta = 2.5104(2)$



$T/T_c = 0.9925; L = 6.0 \text{ fm}; 72^3 \times 8$




$T/T_c = 1.0024$; $L = 5.8$ fm; $72^3 \times 8$





Scatter plot: asymmetry versus Polyakov loop; $\tau = 0.008, L = 5.9 \text{ fm}$


Regression analysis




The problem: to find the conditional expectation value of Y as a function of X : $E(Y|X) = f(X, \theta)$, here $f(X, \theta) = \theta_0 + \theta_1 X$
We employ the method of least squares to determine θ_i - the parameters of regression $D(Y|X)$ is needed!










Infinite-volume limit



Find the regression curve

$$\Delta_{A^2} = \Delta_{A^2}^C + A_1^+ \mathcal{P} + A_2^+ \mathcal{P}^2 \quad \mathcal{P} > 0 \quad (11)$$

$$\Delta_{A^2} = \Delta_{A^2}^C + A_1^- \mathcal{P} + A_2^- \mathcal{P}^2 \quad \mathcal{P} < 0$$

$$A_1^+ = A_1^- = A_1, \quad A_2^+ = A_2^- = A_2$$

$$\begin{aligned}\beta_A &= \beta = 0.326419(3), \\ B_A &= A_1 B = -54.02(24)\end{aligned}$$

In the infinite-volume limit distributions tend to dots on the regression curve

$$\Delta_{A^2} = \Delta_{A^2}^C + A_1 \mathcal{P} + A_2 \mathcal{P}^2 + \dots$$

From this expansion it follows that

$$\mathcal{A} = \Delta_{A^2} - \Delta_{A^2}^C \simeq A_1 \mathcal{P} \simeq A_1 B \tau^\beta , \quad (12)$$


whereas, by definition,


$$\mathcal{A} \simeq B_A \tau^{\beta_A} . \quad (13)$$


Therefore,


$$\begin{aligned}\beta_A &= \beta = 0.326419(3), \\ B_A &= A_1 B = -54.02(24)\end{aligned}$$


At $\tau < 0$ $\mathcal{A} \approx 0$ is a smooth function












Homoscedasticity is severely broken



Critical behavior of the bare longitudinal propagator

$$D_L(0) = D_L^C(0) + D_1 \mathcal{P} + D_2 \mathcal{P}^2 + \dots$$

From this expansion it follows that

$$\mathcal{D} = D_L(0) - D_L^C(0) \simeq D_1 \mathcal{P} \simeq D_1 B \tau^\beta , \quad (14)$$


whereas, by definition,


$$\mathcal{D} \simeq \mathcal{D} \tau^{\beta_D}. \quad (15)$$

Therefore,


$$\begin{aligned} \beta_D &= \beta = 0.326419(3), \\ B_D &= D_1 B = -1832(60) \text{ GeV}^{-2} \end{aligned}$$

$$a^{-1} \sim 2.5 \text{ GeV}$$






$0.035 < \mathcal{P} < 0.040$; $L = 4 \text{ fm}$; $1 \rightarrow \tau = 0.002$; $2 \rightarrow \tau = 0.025$



$-0.030 < \mathcal{P} < -0.025$; $L = 6$ fm; $1 \rightarrow \tau = -0.0045$; $2 \rightarrow \tau = 0.0148$



$$L = 6 \text{ fm};$$

$$\tau = 0.0048;$$

$$1 \rightarrow 0.035 < \mathcal{P} < 0.040; 2 \rightarrow -0.030 < \mathcal{P} < -0.025;$$

Conclusions

- ▶ Both the asymmetry and the longitudinal propagator have a significant correlation with the Polyakov loop.
- ▶ Regression analysis reveals the dependence of each of these quantities on the Polyakov loop \mathcal{P} as follows:

$$D \simeq D_0 + D_1 \mathcal{P} + D_2 \mathcal{P}^2$$

- ▶ Such dependence implies that in the infinite-volume limit both Δ_{A^2} and $D_L(0)$

$$\beta_A = \beta_D = \beta = 0.326419(3)$$

- ▶ $B_A = -54.02(24)$, $B_D = -1832(60)$ GeV $^{-2}$ (bare quantities)
- ▶ Scaling in the conditional distribution of $D_L(0)$ is observed